

Nonlinear Luenberger-like observers for some anthracnose models

David Jaures FOTSA MBOGNE^{a*}, Duplex Elvis HOUPA DANGA^b, David BEKOLLE^{b†}

^aDepartment of Mathematics and Computer Science, ENSAI, The University of Ngaoundere

^bDepartment of Mathematics and Computer Science, Fac.Sci., The University of Ngaoundere

Abstract

In this paper, we propose observers for the dynamics of anthracnose disease described in [12]. Spatial and non spatial versions of the observers are given following an approach similar to [12]. The models given in [12] are improved in order to be more realistic. The conditions on parameters are more general. There are also changes in the equations modelling the dynamics of the berry volume (v) and the rot volume (v_r). We study theoretically the proposed observers in terms of well-posedness and convergence. We make several simulations to assess the effectiveness of observers which definitely display fairly good behaviour.

KeyWords— Anthracnose modelling, Nonlinear observer, Stability.

AMS Classification— 93C15, 93C20, 93B07, 93D15.

1 Introduction

Anthracnose is a phytopathology which occurs on several commercial tropical crops. The coffee is concerned by that disease due to the *Colletotrichum kahawae* which is an ascomycete fungus [2, 5, 6, 14, 17, 21, 24]. Several models of the anthracnose dynamics have been proposed in the literature [9, 10, 11, 16, 17, 18, 19, 20, 24] in order to better understand and control it. Recently, in [12, 13] the authors studied the optimal control of a diffusion model of anthracnose with continuous and impulsive strategies. Optimal controls were computed with respect to given cost functionals. The general model proposed in [12] was given by the following equations

$$\partial_t \theta = \alpha(t, x) (1 - w(t, x) \theta) + \operatorname{div} (A(t, x) \nabla \theta), \text{ on } \mathbb{R}_+^* \times U \quad (1)$$

$$\langle A(t, x) \nabla \theta(t, x), n(x) \rangle = 0, \text{ on } \mathbb{R}_+^* \times \partial U \quad (2)$$

where $n(x)$ denotes the normal vector on the boundary at x and

$$\theta(0, x) = \rho(x) \in [0, 1[, \quad x \in \overline{U} \subseteq \mathbb{R}^3 \quad (3)$$

$$\partial_t v = \beta(t, x, \theta) (1 - v / ((1 - \theta) \eta(t, x) v_{\max})) \quad (4)$$

$$\partial_t v_r = \gamma(t, x, \theta) (1 - v_r / v) \quad (5)$$

$$(v(0, x), v_r(0, x)) \in]0, v_{\max}] \times [0, v_{\max}], \quad x \in \overline{U} \subseteq \mathbb{R}^3 \quad (6)$$

where $w(t, x) = 1 / (1 - \sigma u(t, x))$ and $v(0) \geq v_r(0)$.

In the model above, θ is the inhibition rate, v and v_r are respectively the fruit volume density and the rot volume density. The function v is upper bounded by a value $v_{\max}$ which models the natural fact that the fruit growth is limited. Nonnegative functions $\alpha, \beta, \gamma : \mathbb{R}_+ \times U \times \mathbb{R} \rightarrow \mathbb{R}_+$ characterize the effects of environmental and climatic conditions on the rate of change of inhibition rate, fruit volume, and infected

*Corresponding author Address: email: mjdavidfotsa@gmail.com, P.O. Box 455, ENSAI, The University of Ngaoundere

†Co-authors emails: e_houpa@yahoo.com, bekolle@yahoo.fr

fruit volume respectively [9, 10, 11]. There is a control u representing the chemical strategy consisting on the effects after application of fungicides. The positive term $1 - \sigma \in]0, 1[$ models the inhibition rate corresponding to epidermis penetration by hyphae. Once the epidermis has been penetrated, the inhibition rate cannot fall below $1 - \sigma$, even under the maximum control effort ($u = 1$). In the absence of control effort ($u = 0$), the inhibition rate should increase towards 1. The effects of environmental and climatic conditions on the maximum fruit volume are represented by the parameter η which is a $]0, 1[$ -valued function. There is a spatial spread of the disease in the open domain $U \subset \mathbb{R}^3$ of class C^1 through the diffusion term $\operatorname{div}(A \nabla \theta)$. The boundary condition $\langle A \nabla \theta, n \rangle = 0$ where A is a 3×3 -matrix (a_{ij}) could be viewed as the law steering migration of the disease between U and its exterior. For instance, if A is reduced to the identity matrix I then $\langle \nabla \theta, n \rangle = 0$ means that the domain U does not have any exchange with its exterior.

Also in [12] a within-host model has been studied and given by the following equations :

$$d_t \theta = \alpha(t) (1 - w(t) \theta) \quad (7)$$

$$d_t v = \beta(t, \theta) (1 - v/\eta(t) v_{\max} (1 - \theta)) \quad (8)$$

$$d_t v_r = \gamma(t, \theta) (1 - v_r/v) \quad (9)$$

$$(\theta(0), v(0), v_r(0)) \in [0, 1[\times]0, v_{\max}] \times [0, v_{\max}] \quad (10)$$

where $w(t) = 1/(1 - \sigma u(t))$ and $v(0) \geq v_r(0)$. The parameters in (7) – (10) still have the same interpretations given in the general model (1) – (6).

In several cases, especially for the results in [12, 13] on anthracnose disease, an optimal control strategy is given as a feedback and needs to know the state of the system and parameters values. It could be difficult to know exactly the dynamics of the inhibition rate while it is easier to observe volumes v and v_r . The aim of this paper is to provide some observers for the state θ in the models (1) – (6) and (7) – (10) assuming that volumes v and v_r are known. An observer¹ is a function $\hat{\theta}$ having its own dynamics depending on the available observations (v and v_r) and converging towards a neighborhood of θ . Let us recall the following useful adapted definitions about the stability of an observer. The authors refers to [8, 15, 22, 23] and references therein for more general and classical definitions.

Definition 1.1 Assume that θ is S -valued. An observer $\hat{\theta}$ is said to be

- (i) (Locally) stable if there are two neighborhoods $U, V \subseteq S$ of 0 and a time $t_0 > 0$ such that if $\theta(0) - \hat{\theta}(0) \in U$ then $\forall t > t_0, \theta(t) - \hat{\theta}(t) \in V$;
- (ii) (Locally) asymptotically stable if there is a neighborhood $U \subseteq S$ of 0 such that if $\theta(0) - \hat{\theta}(0) \in U$ then $\lim_{t \rightarrow \infty} \theta(t) - \hat{\theta}(t) = 0$;
- (iii) (Locally) exponentially asymptotically stable if there are a neighborhood $U \subseteq S$ of 0, a positive real constant k and a time $t_0 > 0$ such $\forall t > t_0, \|\theta(t) - \hat{\theta}(t)\| < \exp(-kt)$.

Those local stability properties are said to be global if one can extend the set U to the whole set S .

The remainder of the work is organized as it follows. In section 2 we present some new realistic considerations made for the models. We also show that the new model is well-posed. We design observers for the non spatial model in section 3. A theoretical study is done in subsection 3.1 and we make numerical evaluations in subsection 3.2. We also design observers for the spatial model in section 4 and we study them theoretically in subsection 4.1. A numerical evaluation is carried out in subsection 4.2. Finally, there is a conclusion in section 5.

¹See [22] for more details on the topic.

2 New modelling of anthracnose dynamics

In this section we introduce some changes in the models given in [12] based on some remarks in order to become more realistic. Our modifications are especially made on the description of the dynamics of v and v_r .

Looking at equation (4) we can see that when θ tends to the value 1, v tends with an infinite speed to zero. However, we think that either there is a disease or not the berries will reach and remain greater than a minimal volume. Indeed, the disease can not be triggered when the berry is less than a minimal size as for instance the bud size. Even if the fruit totally rots it keeps a minimal size. We then suggest to add a very small nonnegative term $\varepsilon \ll 1$ in such way that equation (4) becomes

$$\partial_t v = \beta(t, x, \theta) (1 - v / ((1 + \varepsilon - \theta) \eta(t, x) v_{\max})) \quad (11)$$

With the same arguments equation (8) becomes

$$d_t v = \beta(t, \theta) (1 - v / \eta(t) v_{\max} (1 + \varepsilon - \theta)) \quad (12)$$

The use of the term ε generalizes the model in such way that we can recover the original model given in [12] by setting $\varepsilon = 0$. With a positive ε even if $\theta = 1$, v might be also positive. Indeed, the berries reach a minimal volume in the neighborhood of $\varepsilon v_{\max} \min_t \{\eta(t)\}$.

We now focus on the equations (5) and (9). In [12] it was established that v_r is $[0, v_{\max}]$ valued. However, that behaviour is not sufficient. Indeed, v_r should remain less than v . On the other hand v_r should remain equal to zero while v is less than the minimal value corresponding to the threshold volume where the disease can be triggered. We then suggest that v_r remains equal to zero when v is not greater than 0. Instead of studying directly v_r , we introduce a proportion function ρ depending on time and space variables, and given such as

$$v_r = \rho v \quad (13)$$

We propose the following dynamics for the rot proportion ρ : $\forall x \in \overline{U}$, $\rho(0, x) \in [0, 1]$ and

$$\partial_t \rho(t, x) = \overline{\gamma}(t, x, \theta(t, x), v(t, x), \rho(t, x)) (1 - \rho(t, x)), \quad \forall (t, x) \in \mathbb{R}_+^* \times \overline{U}, \quad (14)$$

where $\overline{\gamma} : \mathbb{R}_+ \times U \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function satisfying $\forall (t, x) \in \mathbb{R}_+ \times U$,

Assumption 2.1 $\overline{\gamma}$ is a measurable with respect to the two first parameters and locally Lipschitz continuous with respect to the third parameter.

Assumption 2.2 $\overline{\gamma}(t, x, \cdot, \cdot, 0)$ is nonnegative and such that

$$\overline{\gamma}(t, x, 0, \cdot, 0) = 0 = \overline{\gamma}(t, x, \cdot, 0, \cdot). \quad (15)$$

Assumption 2.3 $\overline{\gamma}(t, x, \cdot, \cdot, \cdot)$ is increasing with respect to the first parameter and $\overline{\gamma}(t, x, 0, \cdot, \cdot)$ is decreasing with respect to the last parameter.

For instance $\overline{\gamma}$ could be chosen with the simple but general form $\overline{\gamma}(t, x, y_1, y_2, y_3) = (\gamma_1(t, x) y_1 - \gamma_2(t, x) y_3) y_2$, with $\gamma_1, \gamma_2 : \mathbb{R}_+ \times U \times \mathbb{R}^3 \rightarrow \mathbb{R}_+$. If we omit the dependence with the space variable we get the corresponding equation for the non spatial model: $\rho(0) \in [0, 1]$ and

$$d_t \rho(t) = \overline{\gamma}(t, \theta(t), v(t), \rho(t)) (1 - \rho(t)), \quad \forall t \in \mathbb{R}_+^*. \quad (16)$$

As we will check in the sequel, the function ρ is $[0, 1]$ -valued. The assumption (2.2) guarantees that while the berry has a null volume (without berry) or the disease has not started the rot volume remains null. The assumption (2.3) means that the rot volume increases with the inhibition rate. When there is a not inhibition the volume of rot does not increase while the fruit grows better and therefore the proportion ρ decreases.

2.1 Well-posedness of the problem for the non spatial model

This subsection is devoted to the proof of the well-posedness of the non spatial model. We make the following useful assumptions.

Assumption 2.4 $\alpha \in L^\infty(\mathbb{R}_+; [0, 1])$.

Assumption 2.5 $u, \eta \in L^\infty(\mathbb{R}_+; [0, 1/(1 + \varepsilon)])$ and $\forall t \geq 0, \eta(t) \geq \eta^* \in]0, 1[$.

Assumption 2.6 $\beta : \mathbb{R}_+ \times U \times \mathbb{R} \rightarrow \mathbb{R}_+$ is nonincreasing function with respect to the third parameter. $\forall z \in \mathbb{R}, \beta(., ., z)$ is a measurable function and belongs to the space $L_{loc}^\infty(\mathbb{R}_+ \times \mathbb{R}; \mathbb{R})$.

Assumption 2.7 $\forall t \geq 0$, the function $\beta(t, .)$ is differentiable and $\partial_x \beta(t, .) \in L_{loc}^\infty(\mathbb{R}; \mathbb{R})$.

The assumptions (2.4)-(2.6) are more general than those given in [12] in the sense that all the parameters of the model were assumed continuous with respect to the state variable while here they are just measurable and essentially bounded. The following existence and uniqueness proposition holds:

Proposition 2.1 If $(\theta(0), v(0), \rho(0)) \in [0, 1] \times [0, v_{\max}] \times [0, 1]$ then the problem (7), (12) and (16) has a unique absolute continuous solution (θ, v, ρ) valued in the set $[0, 1] \times [0, v_{\max}] \times [0, 1]$.

Proof. Using the Carathéodory theorem (see [7]) we get easily the existence of an absolutely continuous solution. The logistic form of the equations (7) and (12) shows clearly that for a maximal solution (θ, v, v_r) the couple (θ, v) is necessarily valued in the set $[0, 1] \times]0, v_{\max}]$. On the other hand from the equation (16), ρ is valued in $[0, 1]$. To show it let set $\rho_1 = \max\{0, -\rho\}$ and $\rho_2 = \max\{0, \rho - 1\}$. The functions ρ_1 and ρ_2 are continuous as ρ is continuous. Without loss of generality we will assume that ρ_1 and ρ_2 are positive on an open interval of time $]0, T[$. Then $\forall t \in]0, T[$,

$$\begin{aligned} \rho_1(t) &= - \int_0^t \overline{\gamma}(s, \theta(s), v(s), -\rho_1(s)) (1 + \rho_1(s)) ds \\ &\leq 0 \end{aligned} \tag{17}$$

$$\begin{aligned} \rho_2(t) &= - \int_0^t \overline{\gamma}(s, \theta(s), v(s), \rho_2(s) + 1) \rho_2(s) ds \\ &\leq 0 \end{aligned} \tag{18}$$

There is a contradiction meaning that ρ_1 and ρ_2 are identically null. Therefore, ρ is $[0, 1]$ -valued. ■

2.2 Well-posedness of the problem for the spatial model

In this subsection the well-posedness of the spatial model is proved. To that aim, we make some assumptions.

Assumption 2.8 $\alpha \in L_{loc}^\infty(\mathbb{R}_+; L^\infty(U; \mathbb{R}_+))$.

Assumption 2.9 $u, \eta \in L^\infty(\mathbb{R}_+; L^\infty(U; [0, 1]))$ and $\forall t \geq 0, 1/(1 + \varepsilon) \geq \eta(t, .) \geq \eta^* \in L^\infty(U; [0, 1])$ with $0 < \inf_U(\eta^*) \stackrel{\text{def}}{=} \eta_m^*$.

Assumption 2.10 $\beta : \mathbb{R}_+ \times U \times \mathbb{R} \rightarrow \mathbb{R}_+$ is nonincreasing function with respect to the third parameter. $\forall z \in \mathbb{R}, \beta(., ., z)$ is a measurable function and belongs to the space $L_{loc}^\infty(\mathbb{R}_+ \times \mathbb{R}; \mathbb{R}_+)$.

Assumption 2.11 $\forall i, j \in \{1, 2, 3\}, a_{ij} \in L_{loc}^\infty(\mathbb{R}_+; W^{1, \infty}(U; \mathbb{R}))$.

Assumption 2.12 $\exists \zeta \in \mathbb{R}_+^*$ such that $\forall t \in \mathbb{R}_+, \forall w \in H^1(U; \mathbb{R})$,

$$\int_U \langle A(t, x) \nabla w(x), \nabla w(x) \rangle dx \geq \zeta \int_U \langle \nabla w(x), \nabla w(x) \rangle dx.$$

Assumption 2.13 $\forall t \geq 0, \forall x \in U$, the function $\beta(t, x, \cdot)$ is differentiable and $\partial_y \beta(t, x, \cdot) \in L_{loc}^\infty(\mathbb{R}; \mathbb{R})$.

Assumptions (2.8)-(2.10) are also more general than those given in [12] in the sense that all the parameters of the model were assumed continuous with respect to the state variable while here they are just measurable and essentially bounded. From assumptions (2.11)-(2.12) the following problem has a unique solution in $H^1(U)$ for an arbitrary but fixed time $t > 0$.

$$\begin{cases} \operatorname{div}(A(t, x) \nabla w(x)) = f(x), \quad \forall x \in U \\ \langle A(t, x) \nabla w(x), n(x) \rangle = 0, \quad \forall x \in \partial U \end{cases}$$

where $f \in L^2(U)$. Following [3] in Theorems 3.6.1 and 3.6.2, there is an orthonormal complete system $\{\varphi_n(t, \cdot)\}_{n \in \mathbb{N}} \subset L^2(U)$ of eigenfunctions and eigenvalues $\{\lambda_n(t)\}$ such that $\forall n \in \mathbb{N}$,

$$\begin{cases} \operatorname{div}(A(t, x) \nabla \varphi_n(t, x)) = \lambda_n(t) \varphi_n(t, x), \quad \forall x \in U \\ \langle A(t, x) \nabla \varphi_n(t, x), n(x) \rangle = 0, \quad \forall x \in \partial U \end{cases}.$$

Moreover, $\{\varphi_n(t, \cdot)\}_{n \in \mathbb{N}} \subset H^1(U)$ and if ∂U is of class C^2 then $(\varphi_n(t, \cdot))_{n \in \mathbb{N}}$ is $H^2(U)$ -valued.

Now we make the following additional assumption.

Assumption 2.14 The sequence (φ_n) does not depends on the time (ie $\varphi_n(t, \cdot) = \varphi_n(\cdot)$, $\forall t > 0$).

Assumption (2.14) could happen if $A(t, \cdot)$ has the form $\mu(t) B(\cdot)$ with $\mu(t) \in \mathbb{R}$, $\forall t \geq 0$. It will be the case in particular if $A(t, \cdot) = \mu(t) I$ and therefore $\operatorname{div}(A(t, \cdot) \nabla w) = \mu(t) \Delta w$. Here I denotes the identity matrix of $\mathbb{R}^3$. Whether (2.14) is satisfied a weak solution θ of (1) – (3) can be written as the sum $\sum_{n=0}^\infty \theta_n \varphi_n$ where each θ_n is an absolutely continuous function of the time and satisfies $\forall t \geq 0$,

$$\begin{aligned} \theta_n(t) &= \sum_{m=0}^\infty \theta_m \int_U \varphi_m(x) \varphi_n(x) dx \\ &= \theta_n(0) + \int_0^t \lambda_n(s) \theta_n(s) ds + \int_0^t \int_U \alpha(s, x) \varphi_n(x) ds dx \\ &\quad - \int_0^t \int_U \alpha(s, x) w(s, x) \theta(s, x) \varphi_n(x) dx ds. \end{aligned} \tag{19}$$

Proposition 2.2 The problem (1) – (3) has a unique solution $\theta \in C(\mathbb{R}_+; H^1(U; [0, 1]))$ which is absolutely continuous with respect to the time.

Proof. It is sufficient here to establish that if there is a maximal solution then that solution is valued in the set $[0, 1]$. Indeed, conditions (2.8)-(2.12) are sufficient to use the Carathéodory theorem (see in [1] the methods used in the proof of the Lemma A 2.7, p187-191). Let $\forall t \geq 0$, $\theta_1 = \max\{0, -\theta\}$ and $\theta_2 = \max\{0, \theta - 1\}$. We have $\theta_1(0, \cdot) = \theta_2(0, \cdot) = 0$. Let S_i be a convex open subset of $\mathbb{R}_+$ where functions $\theta_i(t, \cdot)$ ($i \in \{1, 2\}$) are positive on subsets of U with a positive measure. Then $\forall t \in S_i$ we have

$$\begin{aligned} \partial_t \|f_1(t, \cdot)\|_{L^2(U; \mathbb{R})}^2 / 2 &= \int_U f_1(t, x) \partial_t f_1(t, x) dx \\ &= - \int_U \alpha(t, x) (1 + w(t, x) f_1(t, x)) f_1(t, x) dx \\ &\leq 0 \end{aligned}$$

$$\begin{aligned}
\partial_t \|f_2(t, \cdot)\|_{L^2(U; \mathbb{R})}^2 / 2 &= \int_U f_2(t, x) \partial_t f_2(t, x) dx \\
&= \int_U \alpha(t, x) (1 - w(t, x)) f_2(t, x) dx - \int_U \alpha(t, x) w(t, x) f_2^2(t, x) dx \\
&\leq 0
\end{aligned}$$

Using the above inequalities we get that functions $\theta_i(t, \cdot)$ are nonpositive on the sets S_i which are necessary empty. We conclude that $\forall t > 0$, $\theta(t, \cdot)$ is valued in $[0, 1]$. ■

Proposition 2.3 *The problem (11), (14) has a unique solution $(v, \rho) \in C(\mathbb{R}_+; L^\infty(U; [0, v_{\max}] \times [0, 1]))$ which is absolutely continuous with respect to the time.*

Proof. Since θ is valued in the set $[0, 1]$ and there is not diffusion in the equations (11) and (14), we can fix the space variable and use a proof similar to the one given for proposition 2.1. ■

3 Observation for the within-host model

3.1 Theoretical design of observers

The main objective of this subsection is to design observers for the problem (7), (12) and (16) in order to estimate the inhibition rate θ . Let us consider the following system : $\forall t \geq 0$,

$$d_t \hat{\theta}(t) = \alpha(t) (1 - w(t) \hat{\theta}(t)) + k_1(t) \phi_1(t, \hat{\theta}(t), \hat{v}(t)) + k_2(t) \phi_2(t, \hat{\theta}(t)) \quad (20)$$

and

$$d_t \hat{v}(t) = \beta(t, \hat{\theta}(t)) \phi_3(t, \hat{\theta}(t), \hat{v}(t)) \quad (21)$$

where

$$\begin{aligned}
\phi_1(t, x, y) &= \begin{cases} (1 - v(t)/y) (1 + \varepsilon - x), & \text{if } v(t) \leq y \text{ and } (x, y) \in]0, 1[\times \mathbb{R}_+^* \\ 0, & \text{otherwise} \end{cases} \\
\phi_2(t, x) &= \begin{cases} d_t \rho(t) - \bar{\gamma}(t, x, v(t), \rho(t)) (1 - \rho(t)), & \text{if } x \in]0, 1[\\ 0, & \text{otherwise} \end{cases}
\end{aligned}$$

and

$$\phi_3(t, x, y) = 1 - \frac{y}{(1 + \varepsilon - x) \eta(t) v_{\max}}.$$

We make the following additional hypothesis :

Assumption 3.1 $k_1, k_2 \in L_{loc}^\infty(\mathbb{R}_+; \mathbb{R}_+)$.

By a solution of the system (20) – (21) we mean an absolutely continuous function of the time $(\hat{\theta}, \hat{v})$ which satisfies $\forall t > 0$,

$$\begin{aligned}
\hat{\theta}(t) &= \hat{\theta}(0) + \int_0^t \alpha(s) (1 - w(s) \hat{\theta}(s)) + \int_0^t k_1(s) \phi_1(s, \hat{\theta}(s), \hat{v}(s)) ds \\
&\quad + \int_0^t k_2(s) \phi_2(s, \hat{\theta}(s)) ds
\end{aligned}$$

and

$$\hat{v}(t) = \hat{v}(0) + \int_0^t \beta(s, \hat{\theta}(s)) \phi_3(s, \hat{\theta}(s), \hat{v}(s)) ds.$$

We have the existence and uniqueness

Proposition 3.1 *If $(\widehat{\theta}(0), \widehat{v}(0)) \in [0, 1] \times [0, v_{\max}]$ then the problem (20) – (21) has a unique solution valued in $[0, 1] \times [0, v_{\max}]$.*

Proof. We first show that if $(\widehat{\theta}, \widehat{v})$ is a local solution of the system (20) – (21) then $(\widehat{\theta}, \widehat{v})$ is valued in $[0, 1] \times [0, v_{\max}]$. Let $\forall t \geq 0$, $\theta_1 = \max\{0, -\widehat{\theta}\}$, $\theta_2 = \max\{0, \widehat{\theta} - 1\}$, $v_1 = \max\{0, -\widehat{v}\}$ and $v_2 = \max\{0, \widehat{v} - v_{\max}\}$. We have $\theta_i(0) = v_i(0) = 0$, $\forall i \in \{1, 2\}$. Without loss of generality we can assume that the functions θ_i and v_i ($i \in \{1, 2\}$) are positive on an open interval $]0, T[$. Then we have

$$\begin{aligned} \theta_1(t) &= - \int_0^t \alpha(s) (1 + w(s) \theta_1(s)) ds - \int_0^t k_1(s) \phi_1(s, -\theta_1(s), \widehat{v}(s)) ds \\ &\quad - \int_0^t k_2(s) \phi_2(s, -\theta_1(s)) ds \\ &\leq 0 \end{aligned}$$

and

$$\begin{aligned} \theta_2(t) &= \int_0^t \alpha(s) (1 - w(s) - w(s) \theta_2(s)) ds + \int_0^t k_1(s) \phi_1(s, 1 + \theta_2(s), \widehat{v}(s)) ds \\ &\quad + \int_0^t k_2(s) \phi_2(s, 1 + \theta_2(s)) ds \\ &\leq 0. \end{aligned}$$

That is θ_1 and θ_2 are identically null and $\widehat{\theta}$ is $[0, 1]$ -valued. In the same manner one can show that $\widehat{v}$ is valued in $[0, v_{\max}]$.

Now let show the existence and the uniqueness of the solution of (20) – (21). It suffices to establish existence and uniqueness of a local solution and use the fact every local solution of (20) – (21) is bounded to conclude using Theorem 5.7 in [4].

Let consider the function $F : (t, x, y) \in \mathbb{R}_+ \times [0, 1] \times [0, v_{\max}] \mapsto \mathbb{R}^2$ defined by

$$F^1(t, x, y, z) = \alpha(t) (1 - xw(t)) + k_1 \phi_1(t, x, y) + k_2 \phi_2(t, x),$$

and

$$F^2(t, x, y) = \beta(t, x) \phi_3(t, x, y).$$

The function F is measurable with respect to the time t and continuous with respect to (x, y) . Using the Carathéodory theorem there is a local solution of (20) – (21). Moreover, F is Lipschitz continuous with respect to (x, y, z) . Therefore, the solution is unique and global. Note that the hypothesis (2.7) implies that $\forall t \geq 0$, the function $\beta(t, \cdot)$ is Lipschitz continuous and bounded on the set $[0, 1]$. ■

Now we state the main results of this subsection. Let define the function $\delta : \mathbb{R} \rightarrow \{0, 1\}$ such as

$$\delta(x) = \begin{cases} 1, & x \in]0, 1[\\ 0, & \text{otherwise} \end{cases}.$$

Theorem 3.2 *Let consider the system (20) – (21) and assume that $\widehat{v}(0) = v(0)$.*

(i) *If the functions k_1 and k_2 are identically null and*

$$\inf \{\alpha(t); t > 0\} > 0 \tag{22}$$

then $\widehat{\theta}$ is a globally exponentially asymptotically stable observer for θ .

- (ii) If the function k_1 is identically null and k_2 is not identically null then $\hat{\theta}$ is a a locally asymptotically stable observers for θ . Moreover, if there is a positive function $C_\gamma \in L_{loc}^\infty(\mathbb{R}_+; \mathbb{R}_+)$ such that

$$|\overline{\gamma}(t, y_1, y_2, y_3) - \overline{\gamma}(t, z_1, z_2, z_3)| \geq C_\gamma(t) \|y - z\| \quad (23)$$

then $\hat{\theta}$ is a a globally exponentially asymptotically stable observers for θ .

- (iii) If the function k_2 is identically null and $\forall t > 0$,

$$\alpha w + k_1 \delta(\hat{\theta}) + \frac{k_1 \delta(\hat{\theta}) \beta(t, \theta) (\eta v_{\max} (1 + \varepsilon - \theta) - v)}{\alpha \eta v v_{\max} (1 - \theta w)} > 0. \quad (24)$$

then $\hat{\theta}$ is a a locally exponentially asymptotically stable observers for θ .

- (iv) If the functions k_1 and k_2 are not identically null, and $\forall t > 0$,

$$\alpha w + k_1 \delta(\hat{\theta}) + k_2(t) \phi_2(t, \hat{\theta}) + \frac{k_1 \delta(\hat{\theta}) \beta(t, \theta) (\eta v_{\max} (1 + \varepsilon - \theta) - v)}{\alpha \eta v v_{\max} (1 - w \theta)} > 0 \quad (25)$$

then $\hat{\theta}$ is a a locally exponentially asymptotically stable observers for θ . Moreover, if the condition (23) is satisfied and

$$k_2 \left| \phi_2(t, \hat{\theta}) \right| > k_1 \phi_1(t, \hat{\theta}, \hat{v}) \quad (26)$$

then $\hat{\theta}$ is a a globally exponentially asymptotically stable observers for θ .

Proof. Let $\hat{e} = \theta - \hat{\theta}$ be the error of estimation.

- (i) The error $\hat{e}$ satisfies

$$d_t \hat{e} = -\alpha(t) w(t) \hat{e} \quad (27)$$

and

$$\hat{e}(t) = \exp \left(- \int_0^t \alpha(s) w(s) ds \right) \hat{e}(0).$$

The result follows.

- (ii) The error $\hat{e}$ satisfies

$$\begin{aligned} d_t \hat{e}^2 &= -2\alpha(t) w(t) \hat{e}^2(t) - 2k_2(t) \phi_2(t, \hat{\theta}(t)) \hat{e}(t) \\ &= -\alpha(t) w(t) \hat{e}^2(t) - 2k_2(t) (1 - \rho(t)) \left(\overline{\gamma}(t, \theta(t), v(t), \rho(t)) - \overline{\gamma}(t, \hat{\theta}(t), v(t), \rho(t)) \right) \hat{e}(t) \\ &\leq -\alpha(t) w(t) \hat{e}^2(t) \end{aligned} \quad (28)$$

and

$$\hat{e}^2(t) \leq \exp \left(- \int_0^t \alpha(s) w(s) ds \right) \hat{e}^2(0).$$

Moreover, if the condition (23) is satisfied then

$$d_t \hat{e}^2 \leq -\alpha(t) w(t) \hat{e}^2(t) - 2k_2(t) (1 - \rho(t)) C_\gamma(t) \hat{e}^2(t)$$

and

$$\hat{e}^2(t) \leq \exp \left(- \int_0^t (\alpha(s) w(s) + 2k_2(s) (1 - \rho(s)) C_\gamma(s)) ds \right) \hat{e}^2(0).$$

(iii) We have

$$\begin{aligned}\frac{\partial_\theta \beta(s, \theta)}{\beta(s, \theta) - d_t v} &= \frac{d_t v \partial_\theta \beta(s, \theta) - \beta(s, \theta) \partial_\theta d_t v}{(\beta(s, \theta) - d_t v)^2} \\ &= \frac{\beta^2(s, \theta) ((1 + \varepsilon - \theta) \partial_\theta v + v)}{\eta v_{\max} (1 + \varepsilon - \theta)^2 (\beta(s, \theta) - d_t v)^2}\end{aligned}$$

and at the neighborhood of 0,

$$\begin{aligned}\frac{\beta(s, \hat{\theta})}{\beta(s, \hat{\theta}) - d_t \hat{v}} &= \frac{\beta(s, \theta)}{(\beta(s, \theta) - d_t v)} - \frac{\beta^2(s, \theta) ((1 + \varepsilon - \theta) \partial_\theta v + v)}{\eta v_{\max} (1 + \varepsilon - \theta)^2 (\beta(s, \theta) - d_t v)^2} \hat{e} + \hat{e} \mathcal{O}(\hat{e}) \\ &= \frac{\eta v_{\max} (1 + \varepsilon - \theta)}{v} - \frac{\eta v_{\max}}{v} \hat{e} - \frac{\eta v_{\max} (1 + \varepsilon - \theta) \partial_\theta v}{v^2} \hat{e} + \hat{e} \mathcal{O}(\hat{e})\end{aligned}$$

It follows that

$$\begin{aligned}\frac{v}{\hat{v}} &= \frac{v}{\eta v_{\max} (1 + \varepsilon - \theta)} \frac{\beta(s, \hat{\theta})}{\beta(s, \hat{\theta}) - d_t \hat{v}} \\ &= 1 - \left(\frac{1}{1 + \varepsilon - \theta} + \frac{\partial_\theta v}{v} \right) \hat{e} - \frac{v \hat{e}}{\eta v_{\max} (1 + \varepsilon - \theta)} \mathcal{O}(\hat{e}).\end{aligned}$$

finally,

$$\begin{aligned}d_t \hat{e} &= -\alpha(t) w(t) \hat{e} - k_1 \phi_1(t, \hat{\theta}, \hat{v}) \\ &= -\frac{k_1(t) \delta(\hat{\theta}) (1 + \varepsilon - \theta) \partial_\theta v}{v} \hat{e} - \left(\alpha(t) w(t) - k_1(t) \delta(\hat{\theta}) \right) \hat{e} \\ &\quad - \frac{k_1(t) \delta(\hat{\theta}) v}{\eta v_{\max}} \hat{e} \mathcal{O}(\hat{e}) \\ &\approx -\frac{k_1(t) \delta(\hat{\theta}) \beta(t, \theta) (\eta v_{\max} (1 + \varepsilon - \theta) - v)}{\alpha \eta v v_{\max} (1 - \theta w)} \hat{e} - \left(\alpha(t) w(t) + k_1(t) \delta(\hat{\theta}) \right) \hat{e}\end{aligned}$$

The given approximation on the dynamics of $\hat{e}$ shows clearly that $\hat{\theta}$ is a locally exponentially asymptotically stable observers for θ if the condition (24) is satisfied.

(iv) With the same arguments developped in (ii) and (iii) we easily get that $\hat{\theta}$ is a locally exponentially asymptotically stable observers for θ . Using the conditions (23) and (26) we are sure that even the behaviour of the term with ϕ_1 tends to introduce unstability, the stability is maintained by the term with ϕ_2 .

■

3.2 Numerical evaluation of the observers

This section is devoted to numerical simulations in order to check the effectiveness of the observers we studied theoretically in the subsection 3.1. The Theorem 3.2 guarantees asymptotical stability of the observers, but pratically the convergence is needed in a finite time. Since the coffee cultivation campaign is a cyle of one year, our simulations will cover the year. Parameters are taken following [12]. The control strategy u is

taken with many variations in order to study their impact on the performances of the observer. For every time $t \geq 0$, u is given by

$$u(t) = \sin^2 \left(\omega_1 (t - \varphi_1)^2 \right) \exp \left(-\omega_2 (t - \varphi_2)^2 \right). \quad (29)$$

The functions α, β and γ are taken with the following form

$$\alpha(t) = p_1(t) + b_1 (1 - \cos(c_1 t)) (t - d_1)^2, \forall t \in \mathbb{R}_+, \quad (30)$$

$$\beta(t, x) = b_2 (1 - \cos(c_2 t)) (t - d_2)^2 p_2(x), \forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}, \quad (31)$$

and

$$\gamma(t, x_1, x_2, x_3) = b_3 (1 - \cos(c_3 t)) (t - d_3)^2 (x_1 - \kappa x_3) x_2, \forall (t, x) \in \mathbb{R}_+ \times \mathbb{R}^3. \quad (32)$$

p_1 is a nonnegative function of the time t and p_2 is a real positive function of x . $\forall i \in \{1, 2, 3\}$, b_i , c_i , and d_i are positive coefficients corresponding respectively to the maximal amplitude, the pulsation and the global maximum of α, β and γ . κ is a positive constant regulating the evolution of the rot volume with respect to the inhibition rate. The terms $1 - \cos(c_i t)$ represent the seasonality probably due to climatic and environmental variations.

Theoretically the functions k_1 and k_2 might take arbitrary positive numbers and we can set them constant without loss of generality. However, for the stability of the numerical scheme we were constrained to set them less than $1/10\Delta t$. Here Δt denotes the constant time step. The initial conditions are taken such as $\theta(0) \in \{0.05, 0.25, 0.50, 0.75\}$, $v(0) \in \{0, 0.25, 0.50, 0.75\}$, $\rho(0) \in \{0.25, 0.50, 0.75\}$, $\hat{\theta}(0) = 0$, $\hat{v}(0) = v(0)$. Indeed, at the beginning the fruits are small and the inhibition rate is relatively small.

The following table gives the assumed parameters values.

Parameters	Values	Source	Parameters	Values	Source
b_1	$5 \ln(10)$	Assumed	$p_1(t)$	0	Assumed
b_2	$v_{\max} \ln(10^5 v_{\max} (1 - \varepsilon \eta^*)) / 2$	Assumed	$p_2(x)$	$2 - x$ or $(2 - x)^2$	Assumed
b_3	$v_{\max} \ln(10^5 v_{\max})$	Assumed	κ	1	Assumed
d_1	7.5×10^{-1}	[12]	$v_{\max}$	1 cm^2	Assumed
d_2	7.5×10^{-1}	[12]	ε	10^{-4}	Assumed
d_3	7.5×10^{-1}	[12]	$\eta(t)$	$1 / (1 + \varepsilon) \text{ cm}^2$	Assumed
c_1	10π	[12]	σ	0.9	[12]
c_2	10π	[12]	k_1	0 or 10^3	Assumed
c_3	10π	[12]	k_2	0 or 10^3	Assumed
ω_1	25π	Assumed	Δt	10^{-4}	Assumed
ω_2	10	Assumed	φ_1	0.6	Assumed
φ_1	0.4	Assumed			

Table 1: Simulation parameters for the non spatial model

In the remainder of the subsection we present simulations of the observers and their estimation relative errors dynamics. The relative error seems to be a good mean to evaluate the performance. For each scenario we display two groups figures. The first one represents the dynamics both of the inhibition rate and observers corresponding to each values of $\rho(0) \leq \theta(0)$. The second group of figures shows relatives errors of observers corresponding to each values of $\rho(0) \leq \theta(0)$.

3.2.1 Simulations with $\theta(0) = v(0) = 0.05$

The figures 1 and 2 show the behaviour of the observers when the observation is started early. As we can see the global performance of each observer is relatively good but the case $k_1 = 0$ and $k_2 = 10^3$ seems to be the better one.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 1: Inhibition rate and observers for $\theta(0) = v(0) = 0.05$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 2: Relative absolute estimation error for $\theta(0) = v(0) = 0.05$.

3.2.2 Simulations with $\theta(0) = 0.05$ and $v(0) = 0.5$

The figures 3 and 4 correspond to an observation starting at the beginning of the disease but with a well developed berry. We have better performances when $k_2 = 10^3$. With $k_1 = 10^3$ and $k_2 = 0$ we notice an unstability behaviour in figure 4 although the error takes the null value. That may be due to the constant sign of the function ϕ_1 .

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 3: Inhibition rate and observers for $\theta(0) = 0.05$ and $v(0) = 0.5$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 4: Relative absolute estimation error for $\theta(0) = 0.05$ and $v(0) = 0.5$.

3.2.3 Simulations with $\theta(0) = 0.75$ and $v(0) = 0.05$

In the figures 5 and 6 the inhibition rate is already high while the berry is little. As expected the worst estimation corresponds to $k_1 = k_2 = 0$. The best estimations are still made when $k_2 = 10^3$. However, when $k_1 = 10^3$, the same unstability behaviour in figure 4 appears. Note that the performances of the observers are better with $\rho(0)$ small.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 5: Inhibition rate and observers for $\theta(0) = 0.75$ and $v(0) = 0.05$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 6: Relative absolute estimation error for $\theta(0) = 0.75$ and $v(0) = 0.05$.

3.2.4 Simulations with $\theta(0) = 0.75$ and $v(0) = 0.5$

For the figures 7 and 8 the inhibition rate is already high and the fruit is relatively mature when the observation starts. The worst case are still when $k_2 = 0$ and the best are with $k_2 = 10^3$. Again the performances of the observers are better with $\rho(0)$ small. It seems that starting observation with $\theta(0)$ small and $v(0)$ big guarantees better results.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 7: Inhibition rate and observers for $\theta(0) = 0.75$ and $v(0) = 0.5$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 8: Relative absolute estimation error for $\theta(0) = 0.75$ and $v(0) = 0.5$.

4 Observation for the spatial model

The aim of this section is similar to the previous section. We design observers for the spatial dynamical system (1) – (3), (6), (11) and (14) in order to estimate the inhibition rate θ and we improve them using computer simulations.

4.1 Theoretical design of the spatial model observers

Let consider the following systems : $\forall t \geq 0, \forall x \in \overline{U}$,

$$\begin{aligned} \partial_t \widehat{\theta} = & \alpha(t, x) \left(1 - w(t, x) \widehat{\theta} \right) + K_1(t, x) \Phi_1(t, x, \widehat{\theta}, \widehat{v}) \\ & + K_2(t, x) \Phi_2(t, x, \widehat{\theta}) + \operatorname{div} \left(A(t, x) \nabla \widehat{\theta} \right), \text{ on } \mathbb{R}_+^* \times U \end{aligned} \quad (33)$$

$$\left\langle A(t, x) \nabla \widehat{\theta}_1(t, x), n(x) \right\rangle = 0, \text{ on } \mathbb{R}_+^* \times \partial U \quad (34)$$

$$\partial_t \widehat{v} = \beta(t, x, \widehat{\theta}) \Phi_3(t, x, \widehat{\theta}, \widehat{v}) \quad (35)$$

where

$$\Phi_1(t, x, y, z) = \begin{cases} (1 - v(t, x)/z)(1 + \varepsilon - y), & \text{if } v(t, x) \leq y \text{ and } (x, y, z) \in \overline{U} \times]0, 1[\times \mathbb{R}_+^* \\ 0, & \text{otherwise} \end{cases},$$

$$\Phi_2(t, x, y) = \begin{cases} \partial_t \rho(t, x) - \overline{\gamma}(t, y, v(t, x), \rho(t, x))(1 - \rho(t, x)), & \text{if } (x, y) \in \overline{U} \times]0, 1[\\ 0, & \text{otherwise} \end{cases}$$

and

$$\Phi_3(t, x, y, z) = 1 - \frac{z}{(1 + \varepsilon - y) \eta(t, x) v_{\max}}.$$

We make the following assumption:

Assumption 4.1 $K_1, K_2 \in L_{loc}^\infty(\mathbb{R}_+; L^\infty(U; \mathbb{R}_+))$.

By a solution of the system (33) – (35) we mean an absolutely continuous function with respect to the time $(\widehat{\theta}, \widehat{v})$ which satisfies $\forall t > 0, \forall x \in U$,

$$\begin{aligned} \widehat{\theta}(t, x) = & \widehat{\theta}(0, x) + \int_0^t \alpha(s, x) \left(1 - w(s, x) \widehat{\theta}(s, x) \right) ds + \operatorname{div} \left(A(s, x) \nabla \widehat{\theta}_1 \right) ds \\ & + \int_0^t K_1(s, x) \Phi_1(s, x, \widehat{\theta}, \widehat{v}) + \int_0^t K_2(s, x) \Phi_2(s, x, \widehat{\theta}) \\ \widehat{v}(t, x) = & \widehat{v}(0, x) + \int_0^t \beta(s, x, \widehat{\theta}) \Phi_3(s, x, \widehat{\theta}, \widehat{v}) ds \end{aligned}$$

We have the following

Proposition 4.1 *If $(\widehat{\theta}(0, \cdot), \widehat{v}(0, \cdot)) \in L^2(U; [0, 1[\times [0, v_{\max}])$ then the system (33) – (35) has a unique weak solution $(\widehat{\theta}, \widehat{v}) \in C(\mathbb{R}_+; H^1(U; [0, 1])) \times C(\mathbb{R}_+; L^2(U; [0, v_{\max}]))$.*

Proof. We first show that if $(\widehat{\theta}, \widehat{v})$ is a local solution of the system (33) – (35) then $\forall t > 0$, $(\widehat{\theta}(t, \cdot), \widehat{v}(t, \cdot))$ is valued in $[0, 1] \times [0, v_{\max}]$. Let $\forall t \geq 0$, $\widehat{\theta}_1 = \max \{0, -\widehat{\theta}\}$, $\widehat{\theta}_2 = \max \{0, \widehat{\theta} - 1\}$, $\widehat{v}_1 = \max \{0, -\widehat{v}\}$, $\widehat{v}_2 = \max \{0, \widehat{v} - v_{\max}\}$. We have $\widehat{\theta}_1(0, \cdot) = \widehat{\theta}_2(0, \cdot) = \widehat{v}_1(0, \cdot) = \widehat{v}_2(0, \cdot) = 0$. Let S_i^θ and S_i^v be convex open subsets of $\mathbb{R}_+$ where respectively functions $\widehat{\theta}_i(t, \cdot)$ and $\widehat{v}_i(t, \cdot)$ ($i \in \{1, 2\}$) are positive on subsets of U with a positive measure. Then $\forall t \in S_i^\theta$ (respectively S_i^v) we have

$$\begin{aligned} d_t \left\| \widehat{\theta}_1(t, \cdot) \right\|_{L^2(U; \mathbb{R})}^2 / 2 &= \int_U \widehat{\theta}_1(t, x) \partial_t f_1(t, x) dx \\ &= - \int_U \alpha(t, x) \left(1 + w(t, x) \widehat{\theta}_1(t, x) \right) \widehat{\theta}_1(t, x) dx \\ &\quad - \int_U K_1(t, x) \Phi_1 \left(t, x, -\widehat{\theta}_1(t, x), \widehat{v} \right) dx \\ &\quad - \int_U K_2(t, x) \Phi_2 \left(t, x, -\widehat{\theta}_1(t, x) \right) dx \\ &\leq 0 \end{aligned}$$

$$\begin{aligned} d_t \left\| \widehat{\theta}_2(t, \cdot) \right\|_{L^2(U; \mathbb{R})}^2 / 2 &= \int_U \widehat{\theta}_2(t, x) \partial_t \widehat{\theta}_2(t, x) dx \\ &= \int_U \alpha(t, x) (1 - w(t, x)) \widehat{\theta}_2(t, x) dx - \int_U \alpha(t, x) w(t, x) \widehat{\theta}_2^2(t, x) dx \\ &\leq 0 \end{aligned}$$

$$\begin{aligned} \partial_t \left\| \widehat{v}_1(t, \cdot) \right\|_{L^2(U; \mathbb{R})}^2 / 2 &= \int_U \widehat{v}_1(t, x) \partial_t \widehat{v}_1(t, x) dx \\ &= - \int_U \beta(t, x, -\widehat{v}_1(t, x)) \Phi_3 \left(t, x, \widehat{\theta}_1, -\widehat{v}_1(t, x) \right) \widehat{v}_1(t, x) dx \\ &\leq 0 \end{aligned}$$

and

$$\begin{aligned} d_t \left\| \widehat{v}_2(t, \cdot) \right\|_{L^2(U; \mathbb{R})}^2 / 2 &= \int_U \widehat{v}_2(t, x) \partial_t \widehat{v}_2(t, x) dx \\ &\leq \int_U \beta(t, x, \widehat{v}_2(t, x) + v_{\max}) \Phi_3 \left(t, x, \widehat{\theta}_1, \widehat{v}_2(t, x) + v_{\max} \right) \widehat{v}_2(t, x) dx \\ &\leq 0 \end{aligned}$$

Using the above inequalities we get that functions $\widehat{\theta}_i(t, \cdot)$ and $\widehat{v}_i(t, \cdot)$ ($i \in \{1, 2\}$) are nonpositive on the sets S_i^θ and S_i^v which are necessary empty. We conclude that $\forall t > 0$, $(\widehat{\theta}(t, \cdot), \widehat{v}(t, \cdot))$ is valued in $[0, 1] \times [0, v_{\max}]$.

Now let show the existence and the uniqueness of the solution of (33) – (35). It suffices to establish existence of a local solution to conclude the result. Just as decomposition (19) a solution to (33) – (34) can be written as the sum $\sum_{n=0}^{\infty} \widehat{\theta}_n \varphi_n$. Each $\widehat{\theta}_n$ satisfies the reaction

$$\begin{aligned} \widehat{\theta}_n(t) &= \sum_{m=0}^{\infty} \widehat{\theta}_m \int_U \varphi_m(x) \varphi_n(x) dx \\ &= \widehat{\theta}_n(0) + \int_0^t \lambda_n(s) \widehat{\theta}_n(s) ds + \int_0^t \int_U \varphi_n(x) \alpha(s, x) \left(1 - w(s, x) \widehat{\theta}(s, x) \right) dx ds \\ &\quad + \int_0^t \int_U \varphi_n(x) K_1(s, x) \Phi_1 \left(s, x, \widehat{\theta}(s, x), \widehat{v}(s, x) \right) dx ds \\ &\quad + \int_0^t \int_U \varphi_n(x) K_2(s, x) \Phi_2 \left(s, x, \widehat{\theta}(s, x) \right) dx ds \end{aligned}$$

Let $\ell^2(\mathbb{R})$ denote the Hilbert space of real-valued sequences $\{s_n\}_{n \in \mathbb{N}}$ such that $\langle s, s \rangle = \sum_{n \in \mathbb{N}} s_n^2 < \infty$. We set

$$S_\theta = \left\{ s \in \ell^2(\mathbb{R}) ; 0 \leq \sum_{n \in \mathbb{N}} s_n \leq 1 \right\}$$

and

$$S_v = \{ w \in L^2(U; \mathbb{R}) ; \forall x \in U, w(x) \in [0, v_{\max}] \}$$

Let consider the linear operator $\mathcal{G} : D(\mathcal{G}) \rightarrow \ell^2(\mathbb{R}) \times L^2(U; \mathbb{R})$ where $D(\mathcal{G}) \subseteq \mathbb{R}_+ \times \ell^2(\mathbb{R}) \times L^2(U; \mathbb{R})$ defined by

$$\mathcal{G}_n^1(t, y, z) = \lambda_n(t) y_n - \int_U \alpha(t, x) w(t, x) \varphi_n(x) \langle y, \varphi(x) \rangle dx - \int_U K_1(t, x) \varphi_n(x) \delta(\langle y, \varphi(x) \rangle) \langle y, \varphi(x) \rangle dx$$

$$\mathcal{G}^2(t, y, z) = 0$$

$\mathcal{G}$ is the infinitesimal generator of the evolution system E defined such as $\forall n \in \mathbb{N}, \forall t_0^+ \leq s \leq t \leq t_1$,

$$(E^1(s, t)(y, z))_n = \int_U \exp\left(\int_s^t (\lambda_n(\tau) - \alpha(\tau, x) w(\tau, x) - K_1(\tau, x)) d\tau\right) \langle y, \varphi(x) \rangle \varphi(x) dx$$

$$E^2(s, t)(y, z) = z$$

Let also consider the operators $\mathcal{F} : D(\mathcal{F}) \cap (\mathbb{R}_+ \times S_\theta \times S_v) \rightarrow \ell^2(\mathbb{R}) \times L^2(U; \mathbb{R})$ where $D(\mathcal{F}) \subseteq \mathbb{R}_+ \times \ell^2(\mathbb{R}) \times L^2(U; \mathbb{R})$ defined by

$$\begin{aligned} \mathcal{F}_n^1(t, y, z) = & \int_U \varphi_n(x) (\alpha(t, x) + K_2(t, x) \bar{\gamma}(t, x, \theta(t, x), v(t, x), \rho(t, x)) (1 - \rho(t, x))) dx \\ & + \int_U K_1(t, x) \delta(\langle y, \varphi(x) \rangle) \frac{(1 + \varepsilon)(z - v) \varphi_n(x)}{z} dx \\ & + \int_U K_1(t, x) \delta(\langle y, \varphi(x) \rangle) \frac{v \varphi_n(x) \langle y, \varphi(x) \rangle}{z} dx \\ & - \int_U \varphi_n(x) K_2(t, x) \bar{\gamma}(t, x, y, v(t, x), \rho(t, x)) (1 - \rho(t, x)) dx \end{aligned}$$

$$\mathcal{F}^2(t, y, z) = \beta(t, x, \langle y, \varphi(x) \rangle) \Phi_3(t, x, \langle y, \varphi(x) \rangle, z)$$

The function $\mathcal{F}$ is measurable with respect to the time t . We can show as in the proof of proposition 3.1 that $\mathcal{F}$ is Lipschitz continuous with respect to (y, z) and there is a global and unique solution for the system (33) – (35). Note that $\forall (t, x) \in \mathbb{R}_+ \times U$ the function $\beta(t, x, \cdot)$ is Lipschitz continuous and bounded on the set $[0, 1]$ using the assumption (2.13). ■

Theorem 4.2 *Let consider the system (33) – (35) and assume that $\hat{v}(0, \cdot) = v(0, \cdot)$.*

(i) *If the functions K_1 and K_2 are identically null almost everywhere on U and*

$$\inf \{ \alpha(t, x) ; t > 0, x \in U \} > 0 \quad (36)$$

then $\hat{\theta}$ is a globally exponentially asymptotically stable observer for θ .

(ii) *If the function K_1 is identically null almost everywhere on U and K_2 is not identically null at least on a subset of U with positive measure then $\hat{\theta}$ is a a locally asymptotically stable observers for θ . Moreover, if there is a positive function $C_\gamma \in L_{loc}^\infty(\mathbb{R}_+^* \times U; \mathbb{R}_+)$ such that*

$$|\bar{\gamma}(t, x, y_1, y_2, y_3) - \bar{\gamma}(t, x, z_1, z_2, z_3)| \geq C_\gamma(t, x) \|y - z\| \quad (37)$$

then $\hat{\theta}$ is a a globally exponentially asymptotically stable observers for θ .

(iii) If the function K_2 is identically null and

$$\inf \left\{ \frac{v + (1 + \varepsilon - \theta) \partial_\theta v}{v} K_1 \delta(\hat{\theta}_1) + \alpha(t, x) w(t, x); (t, x) \in \mathbb{R}_+ \times U \right\} > 0. \quad (38)$$

then $\hat{\theta}$ is a locally exponentially asymptotically stable observers for θ .

(iv) If the functions K_1 and K_2 are not identically null, and

$$\inf \left\{ K_1 \delta(\hat{\theta}) \frac{v + (1 + \varepsilon - \theta) \partial_\theta v}{v} + K_2 \Phi_2(t, x, \hat{\theta}) + \alpha(t, x) w(t, x); (t, x) \in \mathbb{R}_+ \times U \right\} > 0. \quad (39)$$

then $\hat{\theta}$ is a locally exponentially asymptotically stable observers for θ . Moreover, if the condition (37) is satisfied and

$$K_2(t, x) \left| \Phi_2(t, \hat{\theta}(t, x)) \right| > K_1(t, x) \Phi_1(t, \hat{\theta}(t, x), \hat{v}(t, x)), \forall (t, x) \in \mathbb{R}_+^* \times U \quad (40)$$

then $\hat{\theta}$ is a globally exponentially asymptotically stable observers for θ .

Proof. Let $\hat{e} = \theta - \hat{\theta}$ be the error of estimation.

(i) The error $\hat{e}$ satisfies

$$\begin{aligned} d_t \|\hat{e}(t, \cdot)\|_{L^2(U)}^2 / 2 &= - \int_U \alpha(t, x) w(t, x) \hat{e}^2(t, x) dx \\ &\leq - \|\hat{e}(t, \cdot)\|_{L^2(U)}^2 \inf_{(s, y) \in \mathbb{R}_+ \times U} \{\alpha(s, y)\} \end{aligned} \quad (41)$$

and

$$\|\hat{e}(t, \cdot)\|_{L^2(U)}^2 \leq \exp \left(-2t \inf_{(s, y) \in \mathbb{R}_+ \times U} \{\alpha(s, y)\} \right) \|\hat{e}(0, \cdot)\|_{L^2(U)}^2. \quad (42)$$

We deduce the results.

(ii) The error $\hat{e}$ satisfies

$$d_t \|\hat{e}(t, \cdot)\|_{L^2(U)}^2 / 2 = - \int_U \alpha(t, x) w(t, x) \hat{e}^2 dx - \int_U \langle A(t, x) \nabla \hat{e}, \nabla \hat{e} \rangle dx \quad (43)$$

$$\begin{aligned} &- \int_U K_2(t, x) \Phi_2(t, x, \hat{\theta}) \hat{e}^2(t, x) dx \\ &\leq - \int_U \frac{v + (1 + \varepsilon - \theta) \partial_\theta v}{v} K_1(t, x) \delta(\hat{\theta}) \hat{e}^2 dx - \int_U \alpha(t, x) w(t, x) \hat{e}^2 dx \\ &- \int_U K_2(t, x) (1 - \rho) \left(\bar{\gamma}(t, x, \theta, v, \rho) - \bar{\gamma}(t, x, \hat{\theta}, v, \rho) \right) \hat{e}(t, x) dx \end{aligned} \quad (44)$$

and

$$\|\hat{e}(t, \cdot)\|_{L^2(U)}^2 \leq \exp \left(-2t \inf_{(s, y) \in \mathbb{R}_+ \times U} \{\alpha(s, y)\} \right) \|\hat{e}(0, \cdot)\|_{L^2(U)}^2.$$

Moreover, if the condition (37) is satisfied then

$$\begin{aligned} d_t \|\hat{e}(t, \cdot)\|_{L^2(U)}^2 &\leq - \int_U 2\alpha(t, x) w(t, x) \hat{e}^2(t, x) dx - \int_U 2K_2(t, x) (1 - \rho(t, x)) C_\gamma(t) \hat{e}^2(t, x) dx \\ &\leq -2 \|\hat{e}(t, \cdot)\|_{L^2(U)}^2 \inf_{(s, y) \in \mathbb{R}_+ \times U} \{\alpha(s, y) w(s, y) + K_2(s, y) (1 - \rho(s, y)) C_\gamma(s, y)\} \end{aligned}$$

and

$$\|\hat{e}(t, \cdot)\|_{L^2(U)}^2 \leq \exp \left(-2t \inf_{(s, y) \in \mathbb{R}_+ \times U} \{\alpha(s, y) w(s, y) + K_2(s, y) (1 - \rho(s, y)) C_\gamma(s, y)\} \right) \|\hat{e}(0, \cdot)\|_{L^2(U)}^2.$$

Since the function C_γ is positive the results holds.

(iii) The error $\widehat{e}$ satisfies

$$\begin{aligned}
d_t \|\widehat{e}(t, \cdot)\|_{L^2(U)}^2 / 2 &= - \int_U \alpha(t, x) w(t, x) \widehat{e}^2(t, x) dx - \int_U \langle A(t, x) \nabla \widehat{e}^2(t, x), \nabla \widehat{e}^2(t, x) \rangle dx \\
&\quad - \int_U K_1(t, x) \Phi_1(t, x, \widehat{\theta}_1(t, x), \widehat{v}_1(t, x)) \widehat{e}^2(t, x) dx \\
&\leq - \int_U \frac{v + (1 + \varepsilon - \theta) \partial_\theta v}{v} K_1(t, x) \delta(\widehat{\theta}_1) \widehat{e}^2(t, x) dx \\
&\quad - \int_U \alpha(t, x) w(t, x) \widehat{e}^2(t, x) dx \\
&\approx - \int_U \frac{K_1(t, x) \delta(\widehat{\theta}) \beta(t, \theta) (\eta v_{\max} (1 + \varepsilon - \theta) - v)}{\alpha \eta v v_{\max} (1 - \theta w)} \widehat{e}^2(t, x) dx \\
&\quad - \int_U \left(\alpha(t) w(t) + K_1(t) \delta(\widehat{\theta}) \right) \widehat{e}^2(t, x) dx.
\end{aligned}$$

Using the condition (38) we deduce that $\widehat{\theta}$ is a locally exponentially asymptotically stable observers for θ .

(iv) Using the same arguments developped in (ii) and (iii) we get that $\widehat{\theta}$ is a locally exponentially asymptotically stable observers for θ . With the conditions (37) and (40) we are sure that even if the term with Φ_1 introduces an unstability, the stability is maintained by the term with Φ_2 .

■

4.2 Numerical evaluation of the spatial model's observers

In this section we present numerical simulations in order to check the effectiveness of the observers we studied theoretically in the subsection 4.1. The duration of the simulations is the year. Parameters are taken following [12] and generalize those in the subsection 3.2. We assume a anisotropic radial control strategy u given for every $(t, x) \in \mathbb{R}_+ \times U$,

$$u(t, x) = \sin^2 \left(\|M(x - x_0)\|^2 \right) \sin^2 \left(\omega_1 (t - \varphi_1)^2 \right) \exp \left(-\omega_2 (t - \varphi_2)^2 \right). \quad (45)$$

The matrix M introduces a geometrical anisotropy in the distribution of u . The functions α, β and γ are taken such as

$$\alpha(t, x) = p_1(t, x) + b_1 q_1(x) (1 - \cos(c_1 t)) (t - d_1)^2, \quad (t, x) \in \mathbb{R}_+ \times U \quad (46)$$

$$\beta(t, x, y) = b_2 q_2(x) (1 - \cos(c_2 t)) (t - d_2)^2 p_2(y), \quad (t, x, y) \in \mathbb{R}_+ \times U \times \mathbb{R} \quad (47)$$

and

$$\gamma(t, x, y_1, y_2, y_3) = b_3 q_3(x) (1 - \cos(c_3 t)) (t - d_3)^2 (y_1 - \kappa y_3) y_2, \quad (t, x, y) \in \mathbb{R}_+ \times U \times \mathbb{R}^3. \quad (48)$$

p_1 is a nonnegative function of the time, p_2 is a positive function and for every i in $\{1, 2, 3\}$, q_i is a real anisotropic radial nonnegative function of the space variable as the control u . For every i in $\{1, 2, 3\}$, b_i , c_i , and d_i are positive coefficients corresponding respectively to the maximal amplitude, the pulsation and the global maximum of α, β and γ . For every i in $\{1, 2, 3\}$, q_i is a $[0, 1]$ -valued function of the space variable. The matrix A in equation (1) is assumed to be equal to $10^{-2}I$ because the diffusion is relatively slow. The functions K_1 and K_2 are taken constant and are similar with k_1 and k_2 defined in subsection 3.2. The rest of the parameters are the same described in the table 1. The initial conditions are also taken following the subsection 3.2 and are constant with respect to the space variable in order to easily fulfill the boundary conditions (2). The following table specifies some assumed values of additional parameters: For every $i \in \{1; 2; 3\}$,

Parameters	Values	Source	Parameters	Values	Source
$q_i(x)$	$\frac{\sin^2(\ M_i(x-x_i)\ ^2)+1}{2}$	Assumed	M, M_i	randomly generated	
x_i	0	Assumed		$5 * \text{rand}(3, 3)$	Assumed

Table 2: Simulation parameters for the spatial model

In the remainder of the subsection we present simulations of the observers and their estimation absolute relative error dynamics. To keep the representation simple we just plot the spatial minimum values, the spatial average values and the spatial maximum values. Instead of the estimation relative error we consider the estimation absolute relative error to avoid compensations when averaging. The initial condition $\rho(0, \cdot)$ is always taken equal to $\theta(0, \cdot)$. Indeed, following simulations in subsection 3.2 the performances of the observers decrease with respect to $\rho(0, \cdot)$ and on the other hand, it seems realistic to take $\rho(0, \cdot)$ less than $\theta(0, \cdot)$.

4.2.1 Simulations with $\theta(0, \cdot) = v(0, \cdot) = 0.05$

The figures 9 and 10 show the behaviour of the observers when the observation is started early. As we can see the global performance of each observer is relatively good but the case $k_1 = 0$ and $k_2 = 10^3$ seems to be the better one.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 9: Inhibition rate and observers for $\theta(0, \cdot) = v(0, \cdot) = 0.05$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 10: Relative absolute estimation error for $\theta(0, \cdot) = v(0, \cdot) = 0.05$.

4.2.2 Simulations with $\theta(0, \cdot) = 0.05$ and $v(0, \cdot) = 0.5$

The figures 11 and 12 correspond to an observation starting at the beginning of the disease but with a well developed berry. We have better performances when $k_2 = 10^3$. With $k_1 = 10^3$ and $k_2 = 0$ we notice an unstability behaviour in figure 12 although the error takes the null value. That may be due to the constant sign of the function ϕ_1 .

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 11: Inhibition rate and observers for $\theta(0, \cdot) = 0.05$ and $v(0, \cdot) = 0.5$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 12: Relative absolute estimation error for $\theta(0, \cdot) = 0.05$ and $v(0, \cdot) = 0.5$.

4.2.3 Simulations with $\theta(0, \cdot) = 0.75$ and $v(0, \cdot) = 0.05$

In the figures 13 and 14 the inhibition rate is already high while the berry is little. As expected the worst estimation corresponds to $k_1 = k_2 = 0$. The best estimations are still made when $k_2 = 10^3$. However, when $k_1 = 10^3$, the same unstability behaviour in figure 12 appears.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 13: Inhibition rate and observers for $\theta(0, \cdot) = 0.75$ and $v(0, \cdot) = 0.05$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 14: Relative absolute estimation error for $\theta(0, \cdot) = 0.75$ and $v(0, \cdot) = 0.05$.

4.2.4 Simulations with $\theta(0, \cdot) = 0.75$ and $v(0, \cdot) = 0.5$

For the figures 15 and 16 the inhibition rate is already high and the fruit is relatively mature when the observation starts. The worst case are still when $k_2 = 0$ and the best are with $k_2 = 10^3$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 15: Inhibition rate and observers for $\theta(0, \cdot) = 0.75$ and $v(0, \cdot) = 0.5$.

$$k_1 = k_2 = 0$$

$$k_1 = 0 \text{ and } k_2 = 10^3$$

$$k_1 = 10^3 \text{ and } k_2 = 0$$

$$k_1 = 10^3 \text{ and } k_2 = 10^3$$

Figure 16: Relative absolute estimation error for $\theta(0, \cdot) = 0.75$ and $v(0, \cdot) = 0.5$.

5 Conclusion and prospects

We have studied four observers for the dynamics of anthracnose disease. Our constructions were based on the models given in [12]. We consider two models: a spatial version and a non spatial one stated respectively by the equations (1) – (6) and (7) – (10). Some changes have been made in the original form of those models in order to be more realistic. Precisely, we have changed the equations modelling the dynamics of the berry volume (v) and the rot volume (v_r). Indeed, before the launch of the disease, the berry has to reach a minimal volume. On the other hand the rot volume shall never be greater than total volume. We found that the system describing the disease dynamics can be viewed as the natural observer of the system even the performances are poor. We have studied theoretically two observers which gives fairly good results looking at the relative error. Several simulations have been carried out to assess the effectiveness of our proposed observers. That work has been done for the spatial and the non spatial models.

Although proposed observers seem to display fairly good results, we think that a stochastic approach could be introduced in the estimation process. Indeed, the parameters of the models are submitted to several stochastic variations. Moreover, the measures of v and v_r also contains errors. A mean to take into account those errors could consist in adding noises in our models and proceed to the stochastic filtering by different existing tools. That aspect is studied in on going works of some of the authors of the current paper.

References

- [1] ANITA S., Analysis and Control of Age-Dependent Population Dynamics, Kluwer Academic Publishers, London, 2000.
- [2] BIEYSSE D., BELLA-MANGA D., MOUEN B., NDEUMENI J., ROUSSEL J., FABRE V. and BERRY D., L'antracnose des baies une menace potentielle pour la culture mondiale de l'arabica. plantations, recherche, développement, pp 145-152, 2002.
- [3] BARBU V., Partial Differential Equations and Boundary Value Problems, Kluwer Academic Publishers, Dordrecht, 1998.
- [4] BENZONI-GAVAGE S., Calcul Différentiel et Équations Différentielles, Dunod, Paris, 2010.
- [5] BOISSON C., L'antracnose du caféier, revue de mycologie, 1960.
- [6] CHEN Z., NUNES M., SILVA M., RODRIGUEZ J., Appressorium turgor pressure of *Colletotrichum kahawae* might have a role in coffee cuticle penetration, mycologia, **96**(6), pp. 1199–1208, 2004.
- [7] CODDINGTON E., LEVINSON N., Theory of Ordinary Differential Equations, McGraw-Hill, New York, 1955.
- [8] CORON J. M., Control and Nonlinearity, AMS, Mathematical surveys and Monographs, Vol **136**, 2007.
- [9] DANNEBERGER T., VARGAS J., JONES Jr., and JONES A., A model for weather-based forecasting of anthracnose on annual bluegrass, Phytopathology, Vol. **74**, No. **4**, pp 448-451, 1984.
- [10] DODD J., ESTRADA A., MATCHAM J., JEFFRIES P., JEGER J., The effect of climatic factors on *Colletotrichum gloeosporioides*, causal agent of mango anthracnose, in the Philippines, Plant Pathology (**40**), pp 568-575, 1991.
- [11] DUTHIE J., Models of the response of foliar parasites to the combined effects of temperature and duration of wetness, Phytopathology, Vol. **87**, No. **11**, 1997.
- [12] FOTSA D., HOUPA E., BEKOLLE D., THRON C., NDOUMBE M., Mathematical modelling and optimal control of anthracnose, Biomath, Vol. **3**, 2014.

- [13] FOTSA D., THRON C., Optimal control of anthracnose using mixed strategy, *Math.Biosci.*, Vol. **269**, pp 186-198, 2015.
- [14] JEFFRIES P., DODD J., JEGER M., PLUMBLEY R., The biology and the control of *Colletotrichum* species on tropical fruit crops, *Plant Pathology* (**39**), pp 343-366, 1990.
- [15] LUENBERGER D., *Introduction to Dynamic Systems. Theory, Models, and Applications*, John Wiley & Sons, 1979.
- [16] MOUEN B., BIEYSSE D., CILAS C., and NOTTEGHEM J., Spatio-temporal dynamics of arabica coffee berry disease caused by *Colletotrichum kahawae* on a plot scale. *Plant Dis.* **91**: 1229-1236, 2007.
- [17] MOUEN B., BIEYSSE D., NYASSE S., NOTTEGHEM J., and CILAS C., Role of rainfall in the development of coffee berry disease in *coffea arabica* caused by *Colletotrichum kahawae*, in cameroon, *Plant pathology*, 2009.
- [18] MOUEN B., BIEYSSE D., NJIAYOUOM I., DEUMENI J., CILAS C., and NOTTEGHEM J., Effect of cultural practices on the development of arabica coffee berry disease, caused by *Colletotrichum kahawae*, *Eur J Plant Pathol.* **119**: 391–400, 2007.
- [19] MOUEN B., CHILLET M., JULLIEN A., BELLAIRE L., Le gainage précoce des régimes de bananes améliore la croissance des fruits et leur état sanitaire vis-à-vis de l’anthracnose (*Colletotrichum musae*), *fruits*, vol. **58**, p. 71–81, 2003.
- [20] MOUEN B., NJIAYOUOM I., BIEYSSE D., NDOUMBE N., CILAS C., and NOTTEGHEM J., Effect of shade on arabica coffee berry disease development: Toward an agroforestry system to reduce disease impact, *Phytopathology*, Vol. **98**, No. **12**, 2008.
- [21] MULLER R., L’évolution de l’anthracnose des baies du caféier d’arabie (*coffea arabica*) due à une forme du *Colletotrichum coffeanum* Noack au Cameroun, café cacao thé, vol. **XIV (2)**, 1970.
- [22] TRINH H., FERNANDO T., *Functional Observers for Dynamical Systems*, *Lecture Notes in Control and Information Sciences*, Springer Berlin Heidelberg, 2012.
- [23] TUSCNAK M., G. WEISS, *Observation and Control for Operator Semigroups*, Birkhäuser Verlag AG, Berlin, 2009.
- [24] WHARTON P., DIEGUEZ-URIBEONDO J., The biology of *Colletotrichum acutatum*, *Anales del Jardín Botánico de Madrid* **61(1)**: 3-22, 2004.